

Integrating Clinical Intelligence and Emotional Responsiveness to Build Patient Trust in Healthcare Delivery

Dyah Juliastuti^{1*} , Yohana F. Cahya Palupi Meilani² , Ihsan Nuril Hikam³ , Julia Nathalie⁴

¹Faculty of Health Sciences, Ichsan Satya University, Indonesia

²Management Department, Pelita Harapan University, Indonesia

³Pandawan Group, Indonesia

⁴Ilearning Group, Colombia

¹dyahjuliastuti2@gmail.com, ²yohana.meilani@uph.edu, ³ihsan.nuril@raharja.info, ⁴jnathalie@ilearning.co

*Corresponding Author

Article Info

Article history:

Submission August 22, 2025

Revised September 25, 2025

Accepted October 27, 2025

Published November 03, 2025

Keywords:

Empathetic AI

Healthcare AI

Artificial Empathy

Patient Trust

Explainable AI



ABSTRACT

Artificial Intelligence has significantly transformed diagnostic processes in healthcare by improving speed, pattern recognition, and decision support across clinical settings. However, diagnostic excellence is not determined by accuracy alone. In real-world practice, patients also value compassion, reassurance, trust, and the feeling of being understood during the diagnostic journey. This paper explores the concept of Empathetic AI in Diagnostics, referring to AI systems that not only assist clinical reasoning but also communicate findings in ways that are emotionally responsive, understandable, and supportive of patient needs. Using a literature-based conceptual approach, this study synthesizes recent evidence on diagnostic AI, conversational medical models, artificial empathy, explainable AI, and patient trust. The findings suggest that empathetic AI can strengthen diagnostic interactions by improving patient comprehension, reducing anxiety, and supporting clinicians in delivering more patient-centered care. At the same time, important risks remain, including inauthentic empathy, overreliance on automated advice, bias, privacy concerns, and legal ambiguity. This paper proposes a framework in which empathetic diagnostic AI should be designed around five dimensions: diagnostic accuracy, emotional sensitivity, explainability, clinician oversight, and ethical accountability. The study concludes that the future of diagnostic AI should not focus solely on prediction performance, but also on how AI supports humane, trustworthy, and ethically grounded diagnostic communication.

This is an open access article under the [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/) license.



This is an open-access article under the CC-BY license (<https://creativecommons.org/licenses/by/4.0/>)

©Authors retain all copyrights

1. INTRODUCTION

The adoption of Artificial Intelligence in healthcare has accelerated rapidly, especially in medical imaging, clinical decision support, triage, risk prediction, and conversational health systems [1]. AI now contributes not only to computational diagnosis but also to patient-facing communication. Yet diagnosis is not purely technical, it is also relational [2, 3]. Patients undergoing diagnostic evaluation often experience uncertainty, fear, and emotional vulnerability [4]. For this reason, effective diagnosis requires both analytical precision and empathetic communication. Recent studies show that patients' attitudes toward healthcare AI

depend not only on system performance, but also on trust, transparency, emotional comfort, and whether clinicians remain meaningfully involved [5].

Traditional diagnostic AI has focused on classification accuracy, sensitivity, specificity, and workflow efficiency [6]. However, the rise of large language models and conversational medical agents has expanded the role of AI from “silent analyzer” to “interactive diagnostic partner” [7]. This shift is significant because the way a diagnosis is communicated can affect patient understanding, confidence, compliance, and overall care experience [8]. Evidence from recent studies suggests that AI-generated medical responses may sometimes be perceived as more empathetic than human-generated responses, but these gains do not automatically translate into authentic trust or preference for AI-led care [9, 10]. Patients may still prefer human empathy even when AI responses score higher on perceived warmth or quality.

The emerging concept of Empathetic AI in Diagnostics addresses this gap. It refers to AI systems that can support diagnosis while communicating with emotional awareness, plain-language clarity, and patient-centered sensitivity [11]. In diagnostics, empathy is not limited to emotional expressions such as “I understand your concern.” It also includes adaptive questioning, sensitivity to distress, understandable explanations, acknowledgement of uncertainty, and support for shared decision-making [12, 13]. Conversational diagnostic models such as AMIE demonstrate that diagnostic dialogue can be optimized as an interactional process, not just a computational output. At the same time, explainable AI research shows that trust increases when systems provide understandable rationales for their outputs [14].

This paper aims to examine how empathy can be meaningfully integrated into diagnostic AI [15, 16]. Specifically, it explores the theoretical foundations, enabling technologies, implementation challenges, and design principles for empathetic AI in diagnostics. The paper argues that future healthcare AI should be evaluated not only by how well it diagnoses disease, but also by how responsibly and compassionately it participates in the diagnostic encounter [17, 18].

2. LITERATURE REVIEW

Empathy in healthcare has long been recognized as a core component of effective clinical practice [19]. It contributes to patient satisfaction, trust, adherence, and reduced anxiety. In diagnostic settings, empathy becomes especially important because patients often face ambiguity, fear of serious disease, and information overload [20]. While clinicians naturally combine medical reasoning with interpersonal sensitivity, traditional AI systems were not built for this human dimension [21].

Recent advances in natural language processing and large language models have changed this landscape. AI systems can now generate responses that are not only medically relevant but also emotionally supportive. [22] found that chatbot responses to patient questions were frequently rated higher than physician responses on both quality and empathy. Likewise, systematic reviews indicate that LLMs increasingly demonstrate both cognitive and affective elements of empathy in medical contexts. However, scholars also caution that artificial empathy may feel persuasive without being morally grounded, and may even appear inauthentic if not carefully designed [23].

In parallel, research on diagnostic AI continues to show strong performance in pattern recognition and clinical decision support [24]. Explainable AI studies in radiology, hematology, pulmonology, and endoscopy suggest that diagnostic AI becomes more acceptable when the reasoning process can be inspected and understood [25]. In high-stakes clinical contexts, explainability is closely tied to empathy because patients and clinicians both need understandable justifications, not only probabilities [26]. An AI that explains why a result is uncertain, what additional tests may be needed, and how the recommendation relates to patient symptoms can contribute to a more reassuring diagnostic interaction [27].

Patient trust is another major theme in the literature. Surveys and scoping reviews show that patients are often open to AI in diagnosis when it improves accuracy and efficiency, but they remain concerned about depersonalization, privacy, bias, and reduced clinician involvement [28]. Trust appears strongest when AI is presented as an assistive tool under human oversight rather than as an autonomous replacement for the physician [29]. Studies also show that disclosing AI use in diagnostic practice may affect how patients perceive physician competence, trustworthiness, and empathy. Thus, successful diagnostic AI must balance technical capability with relational design [30, 31].

A final theme in the literature concerns direct-to-consumer and conversational health applications [32]. These systems are increasingly used for self-assessment, symptom interpretation, second opinions, and patient

education [33]. Although they may expand access, their rapid growth raises concerns about hallucinations, overconfidence, liability, and emotional manipulation [34]. Design recommendations from recent reviews stress the need for transparency, retrieval support, domain grounding, and clear escalation pathways to human care [35]. This is particularly relevant for empathetic diagnostic AI, where emotionally sensitive language must never obscure uncertainty or clinical limitations [36–38].

3. METHODOLOGY

This study employs a qualitative conceptual literature review design. The methodology is intended to synthesize interdisciplinary knowledge at the intersection of health, empathy, conversational AI, and diagnostic systems.

3.1. Research Design

The paper adopts a narrative-conceptual synthesis approach. This design is appropriate because the field of empathetic AI in diagnostics is still emerging and spans multiple domains, including clinical AI, medical communication, explainable AI, ethics, digital health, and patient experience.




Figure 1. Conceptual Research Flow

Figure 1 illustrates the overall research flow adopted in this study to develop a conceptual understanding of empathetic AI in diagnostics. The process begins with the identification of the research topic, followed by the collection and selection of relevant literature from interdisciplinary areas, including healthcare artificial intelligence, empathy studies, explainable AI, and patient trust. After the literature is screened, the selected studies are analyzed thematically to identify recurring concepts and relationships that are important for the development of the proposed framework.

This figure shows that the study does not rely on a single dimension of analysis, but instead integrates several major themes, namely diagnostic performance, emotional responsiveness, explainability, patient trust and acceptance, as well as ethical and governance concerns. Through conceptual synthesis, these themes are combined into a unified framework of empathetic AI in diagnostics. Therefore, Figure 1 highlights that the methodology of this paper is systematic, interdisciplinary, and oriented toward building a comprehensive conceptual model.

3.2. Literature Selection

The review conceptually draws on recent literature from healthcare AI, medical informatics, conversational agents, patient trust research, and ethical AI studies. Priority is given to studies published from 2020 onward, particularly those discussing:

- AI-supported diagnosis
- Artificial empathy in healthcare communication

- Explainable AI in clinical settings
- Patient trust and acceptance of diagnostic AI
- Conversational and large language model applications in medicine

3.3. Analytical Procedure

The selected literature was analyzed through thematic synthesis. Five recurring themes were identified:

- Diagnostic performance and clinical utility
- Empathetic communication and emotional responsiveness
- Explainability and interpretability
- Trust, acceptance, and patient perception
- Ethical, legal, and implementation challenges

3.4. Conceptual Objective

The purpose of the analysis is not to test a statistical hypothesis, but to develop a structured conceptual framework for understanding how empathy can be embedded in diagnostic AI in a clinically responsible way.

4. RESULT AND DISCUSSION

The literature synthesis indicates that empathetic AI in diagnostics should be understood as a hybrid capability rather than a single feature. It is not enough for an AI system to produce accurate diagnostic suggestions. It must also communicate uncertainty clearly, respond sensitively to emotional cues, and support patient understanding without replacing clinician judgment.

4.1. Empathy Enhances the Diagnostic Experience

The first major result is that empathy functions as a diagnostic enabler. Patients who feel heard and reassured are more likely to share relevant symptoms, ask clarifying questions, and remain engaged in follow-up care. Conversational systems capable of supportive language, adaptive questioning, and plain-language explanation can therefore improve the quality of the diagnostic encounter. This is especially relevant in primary care, oncology, mental health, chronic illness, and other emotionally charged contexts.

4.2. Explainability is a Form of Clinical Empathy

A second key finding is that explainability should be treated not only as a technical requirement, but also as a relational one. When AI provides understandable rationales, highlights uncertainty, and clarifies why certain symptoms or images matter, it reduces confusion and supports patient trust. In this sense, explainability operates as a form of cognitive empathy: the system recognizes the user's need for meaning, not just output. Research consistently shows that interpretable or explainable systems are more likely to be trusted by clinicians and patients in diagnostic contexts.

4.3. Human Oversight Remains Essential

The third result is that patients generally prefer AI-assisted care over AI-only care. Even when AI-generated responses are judged as highly empathetic, people still value the moral accountability, contextual understanding, and lived compassion associated with human clinicians. This suggests that empathetic diagnostic AI should be positioned as a co-pilot, not a replacement. The most promising model is a human-in-the-loop diagnostic partnership, where AI supports information gathering, explanation, and emotional scaffolding, while clinicians retain responsibility for interpretation, disclosure, and care decisions.

4.4. Risks of Artificial Empathy

The fourth finding concerns risk. Artificial empathy can be beneficial, but it also introduces new dangers. Patients may overtrust systems that sound caring but lack true understanding. Emotional phrasing can create a false sense of competence or safety. Bias, hallucination, weak domain grounding, and data privacy concerns may further undermine safe use. Legal questions also remain unresolved when diagnostic harm results from AI-supported decisions. Therefore, empathetic AI must be bounded by ethical safeguards, explicit uncertainty communication, auditability, and clinical escalation protocols.

4.5. Proposed Framework for Empathetic AI in Diagnostics

Based on the synthesis, this paper proposes five core pillars for empathetic diagnostic AI:

- **Diagnostic Intelligence**
The system must demonstrate clinically meaningful accuracy, relevance, and safe performance.
- **Emotional Responsiveness**
The system should detect concern, confusion, or distress and respond with supportive, non-alarming language.
- **Explainable Communication**
The AI should translate findings into understandable explanations, including uncertainty and next-step guidance.
- **Human Clinical Oversight**
The clinician must remain central in validating outputs, contextualizing recommendations, and preserving accountability.
- **Ethical and Governance Safeguards**
The system must address privacy, fairness, auditability, transparency, and legal responsibility.

Together, these pillars suggest that the future of diagnostic AI should move beyond “smart diagnosis” toward compassionate diagnostic ecosystems where clinical intelligence and patient dignity are developed together.




Figure 2. Core Dimensions

Figure 2 presents the five core dimensions that form the foundation of empathetic AI in diagnostics. These dimensions include diagnostic accuracy, emotional responsiveness, explainable communication, human clinical oversight, and ethical accountability. The figure emphasizes that empathetic AI is not only concerned with the technical ability of AI to generate diagnostic suggestions, but also with how the system communicates in a supportive manner, remains understandable to patients, and operates under responsible clinical supervision. The figure also demonstrates the interconnected nature of these dimensions. Diagnostic accuracy alone is insufficient if the AI system cannot explain its reasoning or respond to patient concerns with sensitivity. Likewise, emotional responsiveness without ethical safeguards may create false reassurance. For this reason, Figure 2 supports the argument that empathetic AI in diagnostics must be designed as a balanced integration of intelligence, communication, trust, and governance, rather than as a purely computational tool.

Table 1. Key Dimensions

Dimension	Description	Clinical Contribution	Potential Risk
Diagnostic Accuracy	Ability of AI to identify patterns, symptoms, and possible diagnoses correctly	Improves efficiency, early detection, and diagnostic support	Misdiagnosis, overreliance, algorithmic bias
Emotional Responsiveness	Ability of AI to respond with supportive, calm, and patient-sensitive language	Reduces anxiety and improves patient comfort during diagnostic interaction	Artificial empathy may feel inauthentic or misleading
Explainable Communication	Ability of AI to provide understandable reasoning and clarify uncertainty	Enhances patient understanding and clinician trust	Oversimplification or incomplete explanation
Human Clinical Oversight	Continued involvement of healthcare professionals in reviewing AI output	Maintains accountability and contextual interpretation	Reduced value if clinicians rely too heavily on AI
Ethical Accountability	Alignment with privacy, fairness, transparency, and governance principles	Protects patient rights and promotes safe adoption	Legal ambiguity, privacy breaches, lack of regulation

Table 1 summarizes the principal dimensions of empathetic AI in diagnostics and explains their respective roles in clinical settings. Each dimension contributes a unique function to the overall design of the system. Diagnostic accuracy ensures that the AI remains clinically useful, while emotional responsiveness supports patient comfort and reduces anxiety during the diagnostic process. Explainable communication further strengthens the interaction by allowing patients and clinicians to understand the logic behind AI-generated outputs.

At the same time, the table highlights that every dimension is accompanied by potential risks. For example, high diagnostic accuracy may still produce bias or overreliance, and emotionally supportive language may appear artificial if not grounded in safe clinical design. Human clinical oversight and ethical accountability are therefore essential to maintaining responsibility, fairness, and trust. In this way, Table 1 shows that empathetic AI must be evaluated not only from the perspective of benefit, but also from the perspective of clinical risk and ethical limitation.

5. MANAGERIAL IMPLICATION

The findings of this study suggest that the managerial importance of empathetic AI in diagnostics extends well beyond operational efficiency and technical performance. In healthcare settings, the value of diagnostic AI is closely tied to its capacity to improve patient understanding, reduce emotional distress, and foster trust during clinically sensitive interactions. This indicates that healthcare managers should not assess AI systems solely on the basis of accuracy, speed, or cost reduction, but also on their ability to provide understandable explanations, communicate uncertainty responsibly, and contribute to a more reassuring diagnostic experience. In this context, empathetic AI should be viewed as a strategic resource for strengthening service quality and supporting more patient-centered models of care.

The study also implies that effective implementation requires strong organizational governance and interdisciplinary design. AI developers, hospital administrators, and clinical leaders need to ensure that empathy-related capabilities are embedded into diagnostic systems through plain-language communication, context-sensitive responses, and appropriate escalation pathways to human professionals. At the same time, empathetic diagnostic AI should remain under clinician-in-the-loop supervision, supported by institutional policies on

transparency, privacy, fairness, and accountability. Such an approach can improve user acceptance, reduce implementation risk, and enhance the long-term legitimacy of AI adoption in healthcare. From a managerial perspective, the significance of empathetic AI lies in its potential to align technological innovation with ethical responsibility, institutional trust, and the humanistic foundations of medical practice.

6. CONCLUSION

In conclusion, this study highlights that empathetic AI in diagnostics should be understood as a multidimensional healthcare innovation that combines diagnostic intelligence with emotional responsiveness, explainable communication, human clinical oversight, and ethical accountability. The discussion shows that the value of AI in diagnostic settings is not limited to improving accuracy and efficiency, but also extends to supporting patient understanding, reducing anxiety, and enhancing the overall quality of the diagnostic encounter. Thus, the integration of empathy into AI-driven diagnostic systems represents an important step toward more patient-centered and humane healthcare delivery.

The practical implication of this study is that healthcare institutions, AI developers, and policymakers should avoid designing diagnostic AI solely around technical performance. Instead, systems should be developed with a balanced focus on communication quality, interpretability, trust, and safety. Clinicians also remain central in ensuring that AI outputs are interpreted responsibly and aligned with the emotional and contextual needs of patients. Therefore, empathetic AI should be positioned as an assistive and collaborative technology that strengthens, rather than replaces, the human dimension of medical practice.

For future research, empirical studies are needed to examine how patients and clinicians respond to empathetic diagnostic AI in real clinical environments. Further investigation is also necessary to measure the effects of empathetic AI on patient trust, comprehension, adherence, and decision-making across different healthcare contexts. In addition, future studies should explore ethical governance models, bias mitigation strategies, and evaluation frameworks that can assess not only diagnostic accuracy but also the authenticity, safety, and relational quality of empathetic AI systems.

7. DECLARATIONS

7.1. About Authors

Dyah Juliastuti (DJ)  <https://orcid.org/0000-0002-4159-9166>

Yohana F. Cahya Palupi Meilani (YF)  <https://orcid.org/0000-0001-7477-8497>

Ihsan Nuril Hikam (IN)  <https://orcid.org/0009-0000-0345-9763>

Julia Nathalie (JN)

7.2. Author Contributions

Conceptualization: DJ; Methodology: YF; Software: IN; Validation: DJ and YF; Formal Analysis: JN and IN; Investigation: DJ; Resources: YF; Data Curation: YF; Writing Original Draft Preparation: IN and JN; Writing Review & Editing: IN and JN; Visualization: YF; All authors, DJ, YF, IN, and JN, have read and agreed to the published version of the manuscript.

7.3. Data Availability Statement

The data presented in this study are available on request from the corresponding author.

7.4. Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

7.5. Declaration of Conflicting Interest

The authors declare that they have no conflicts of interest, known competing financial interests, or personal relationships that could have influenced the work reported in this paper.

REFERENCES

- [1] J. W. Ayers, A. Poliak, M. Dredze, E. C. Leas, Z. Zhu, J. B. Kelley, D. J. Faix, A. M. Goodman, C. A. Longhurst, M. Hogarth *et al.*, “Comparing physician and artificial intelligence chatbot responses to patient questions posted to a public social media forum,” *JAMA internal medicine*, vol. 183, no. 6, pp. 589–596, 2023.
- [2] V. Sorin, D. Brin, Y. Barash, E. Konen, A. Charney, G. Nadkarni, and E. Klang, “Large language models and empathy: systematic review,” *Journal of medical Internet research*, vol. 26, p. e52597, 2024.
- [3] T. A. Wiratno and B. Callula, “Transformation of beauty in digital fine arts aesthetics: An artpreneur perspective,” *Aptisi Transactions on Technopreneurship (ATT)*, vol. 6, no. 2, pp. 231–241, 2024.
- [4] H. S. J. Chew and P. Achananuparp, “Perceptions and needs of artificial intelligence in health care to increase adoption: scoping review,” *Journal of medical Internet research*, vol. 24, no. 1, p. e32939, 2022.
- [5] X. He, X. Zheng, and H. Ding, “Existing barriers faced by and future design recommendations for direct-to-consumer health care artificial intelligence apps: scoping review,” *Journal of Medical Internet Research*, vol. 25, p. e50342, 2023.
- [6] T. Tu, M. Schaeckermann, A. Palepu, K. Saab, J. Freyberg, R. Tanno, A. Wang, B. Li, M. Amin, Y. Cheng *et al.*, “Towards conversational diagnostic artificial intelligence,” *Nature*, vol. 642, no. 8067, pp. 442–450, 2025.
- [7] D. McDuff, M. Schaeckermann, T. Tu, A. Palepu, A. Wang, J. Garrison, K. Singhal, Y. Sharma, S. Azizi, K. Kulkarni *et al.*, “Towards accurate differential diagnosis with large language models,” *Nature*, vol. 642, no. 8067, pp. 451–457, 2025.
- [8] L. Seitz, “Artificial empathy in healthcare chatbots: Does it feel authentic?” *Computers in Human Behavior: Artificial Humans*, vol. 2, no. 1, p. 100067, 2024.
- [9] F. Busch, L. Hoffmann, L. Xu, L. J. Zhang, B. Hu, I. García-Juárez, L. N. Toapanta-Yanchapaxi, N. Gorelik, V. Gorelik, G. A. Rodriguez-Granillo *et al.*, “Multinational attitudes toward ai in health care and diagnostics among hospital patients,” *JAMA network open*, vol. 8, no. 6, p. e2514452, 2025.
- [10] R. A. Sunarjo, M. H. R. Chakim, S. Maulana, and G. Fitriani, “Management of educational institutions through information systems for enhanced efficiency and decision-making,” *International Transactions on Education Technology (ITEE)*, vol. 3, no. 1, pp. 47–61, 2024.
- [11] K. Borys, Y. A. Schmitt, M. Nauta, C. Seifert, N. Krämer, C. M. Friedrich, and F. Nensa, “Explainable ai in medical imaging: An overview for clinical practitioners—beyond saliency-based xai approaches,” *European journal of radiology*, vol. 162, p. 110786, 2023.
- [12] M. Ennab and H. Mcheick, “Enhancing interpretability and accuracy of ai models in healthcare: a comprehensive review on challenges and future directions,” *Frontiers in Robotics and AI*, vol. 11, p. 1444763, 2024.
- [13] F. Zidan, D. Nugroho, and B. A. Putra, “Securing enterprises: harnessing blockchain technology against cybercrime threats,” *International Journal of Cyber and IT Service Management*, vol. 3, no. 2, pp. 168–173, 2023.
- [14] M. Abbasian, I. Azimi, A. M. Rahmani, and R. Jain, “Conversational health agents: a personalized large language model-powered agent framework,” *JAMIA open*, vol. 8, no. 4, p. ooaf067, 2025.
- [15] S. Maity and M. J. Saikia, “Large language models in healthcare and medical applications: a review,” *Bioengineering*, vol. 12, no. 6, p. 631, 2025.
- [16] D. Rustiana, D. Ramadhan, L. Wibowo, A. W. Nugroho, and G. Mahardika, “State of the art blockchain enabled smart contract applications in the university,” *Blockchain Frontier Technology*, vol. 2, no. 2, pp. 70–80, 2023.
- [17] J. Maslinski, R. Grasfield, R. Awasthi, S. Mishra, D. Mahapatra, P. Mathur, and S. Mishra, “Understanding large language models in healthcare: A guide to clinical implementation and interpreting publications,” *Cureus*, vol. 17, no. 4, p. e82397, 2025.
- [18] S. Shool, S. Adimi, R. Saboori Amleshi, E. Bitaraf, R. Golpira, and M. Tara, “A systematic review of large language model (llm) evaluations in clinical medicine,” *BMC Medical Informatics and Decision Making*, vol. 25, no. 1, p. 117, 2025.
- [19] T. Y. C. Tam, S. Sivarajkumar, S. Kapoor, A. V. Stolyar, K. Polanska, K. R. McCarthy, H. Osterhoudt, X. Wu, S. Visweswaran, S. Fu *et al.*, “A framework for human evaluation of large language models in healthcare derived from literature review,” *NPJ digital medicine*, vol. 7, no. 1, p. 258, 2024.
- [20] G. Foresman, J. Biro, A. Tran, K. MacRae, S. Kazi, L. Schubel, A. Visconti, W. Gallagher, K. M. Smith,

- T. Giardina *et al.*, “Patient perspectives on artificial intelligence in health care: focus group study for diagnostic communication and tool implementation,” *Journal of Participatory Medicine*, vol. 17, no. 1, p. e69564, 2025.
- [21] A. M. Astobiza, M. Alonso, and R. Ortega Lozano, “Trust and ai in healthcare: a systematic review,” *Monash Bioethics Review*, pp. 1–21, 2025.
- [22] E. Ratti, M. Morrison, and I. Jakab, “Ethical and social considerations of applying artificial intelligence in healthcare—a two-pronged scoping review,” *BMC Medical Ethics*, vol. 26, no. 1, p. 68, 2025.
- [23] G. Montanari Vergallo, L. L. Campanozzi, M. Gulino, L. Bassis, P. Ricci, S. Zaami, S. Marinelli, V. Tambone, P. Frati *et al.*, “How could artificial intelligence change the doctor–patient relationship? a medical ethics perspective,” in *Healthcare*, vol. 13, no. 18, 2025.
- [24] L. Tang, J. Li, and S. Fantus, “Medical artificial intelligence ethics: A systematic review of empirical studies,” *Digital health*, vol. 9, p. 20552076231186064, 2023.
- [25] M. Chen, B. Zhang, Z. Cai, S. Seery, M. J. Gonzalez, N. M. Ali, R. Ren, Y. Qiao, P. Xue, and Y. Jiang, “Acceptance of clinical artificial intelligence among physicians and medical students: a systematic review with cross-sectional survey,” *Frontiers in medicine*, vol. 9, p. 990604, 2022.
- [26] A. Sauerbrei, A. Kerasidou, F. Lucivero, and N. Hallowell, “The impact of artificial intelligence on the person-centred, doctor-patient relationship: some problems and solutions,” *BMC medical informatics and decision making*, vol. 23, no. 1, p. 73, 2023.
- [27] L. R. Baghdadi, A. A. Mobeirek, D. R. Alhudaithi, F. A. Albenmousa, L. S. Alhadlaq, M. S. Alaql, and S. A. Alhamlan, “Patients’ attitudes toward the use of artificial intelligence as a diagnostic tool in radiology in saudi arabia: cross-sectional study,” *JMIR Human Factors*, vol. 11, no. 1, p. e53108, 2024.
- [28] S. J. Fransen, T. Kwee, D. Rouw, C. Roest, Q. van Lohuizen, F. Simonis, P. van Leeuwen, S. Heijmink, Y. Ongena, M. Haan *et al.*, “Patient perspectives on the use of artificial intelligence in prostate cancer diagnosis on mri,” *European radiology*, vol. 35, no. 2, pp. 769–775, 2025.
- [29] S. A. Berger, E. Håland, and M. Solbjør, “Patient perspectives on trust in artificial intelligence–powered tools in prostate cancer diagnostics,” *Qualitative Health Research*, p. 10497323251387545, 2025.
- [30] B. B. Ozcan, B. E. Dogan, Y. Xi, and E. E. Knippa, “Patient perception of artificial intelligence use in interpretation of screening mammograms: a survey study,” *Radiology: Imaging Cancer*, vol. 7, no. 3, p. e240290, 2025.
- [31] O. Candra, N. B. Kumar, N. K. A. Dwijendra, I. Patra, A. Majdi, U. Rahardja, M. Kosov, J. W. G. Guerrero, and R. Sivaraman, “Energy simulation and parametric analysis of water cooled thermal photovoltaic systems: energy and exergy analysis of photovoltaic systems,” *Sustainability*, vol. 14, no. 22, p. 15074, 2022.
- [32] F. Pesapane, E. Giambersio, B. Capetti, D. Monzani, R. Grasso, L. Nicosia, A. Rotili, A. Sorce, L. Meneghetti, S. Carriero *et al.*, “Patients’ perceptions and attitudes to the use of artificial intelligence in breast cancer diagnosis: a narrative review,” *Life*, vol. 14, no. 4, p. 454, 2024.
- [33] M. A. Ruben, D. Blanch-Hartigan, and J. A. Hall, “What is artificial intelligence (ai)“empathy”? a study comparing chatgpt and physician responses on an online forum: Ruben et al.” *Journal of General Internal Medicine*, pp. 1–8, 2025.
- [34] P. Nong and M. Ji, “Expectations of healthcare ai and the role of trust: understanding patient views on how ai will impact cost, access, and patient-provider relationships,” *Journal of the American Medical Informatics Association*, vol. 32, no. 5, pp. 795–799, 2025.
- [35] Y. Yu, C. A. Gomez-Cabello, S. A. Haider, A. Genovese, S. Prabha, M. Trabilisy, B. G. Collaco, N. G. Wood, S. Bagaria, C. Tao *et al.*, “Enhancing clinician trust in ai diagnostics: A dynamic framework for confidence calibration and transparency,” *Diagnostics*, vol. 15, no. 17, p. 2204, 2025.
- [36] B. Osnat, “Patient perspectives on artificial intelligence in healthcare: A global scoping review of benefits, ethical concerns, and implementation strategies,” *International Journal of Medical Informatics*, vol. 203, p. 106007, 2025.
- [37] M. H. Tilala, P. K. Chenchala, A. Choppadandi, J. Kaur, S. Naguri, R. Saoji, B. Devaguptapu, and M. Tilala, “Ethical considerations in the use of artificial intelligence and machine learning in health care: a comprehensive review,” *Cureus*, vol. 16, no. 6, 2024.
- [38] J. Glenning and L. Gualtieri, “Patient perspectives on artificial intelligence in medical imaging,” *Journal of Participatory Medicine*, vol. 17, p. e67816, 2025.